

Gas-Phase Reactions of X_3M -NCR and
 X_3M -CNR Donor-Acceptor Complexes
($M=Al, Ga; X=H, Cl, CH_3; R=H, CH_3$)

A.Y. Timoshkin, H.F. Schaefer III

published in

Modern Methods and Algorithms of Quantum Chemistry,
J. Grotendorst (Ed.), John von Neumann Institute for Computing,
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 6, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

Gas-phase reactions of $X_3M\text{-NCR}$ and $X_3M\text{-CNR}$ donor-acceptor complexes ($M=\text{Al,Ga}$; $X=\text{H,Cl,CH}_3$; $R=\text{H,CH}_3$).

Alexey Y. Timoshkin^a, Henry F. Schaefer, III^b.

^aSt. Petersburg State University, Department of Chemistry, University pr. 2,
Old Petergof, St.Petersburg, 198904 Russia; e-mail: alex@dux.ru;

^bCenter for Computational Quantum Chemistry, University of Georgia,
Athens, GA 30602, USA. e-mail: hfsiii@arches.uga.edu.

Carbon contamination is one of the major problems during the CVD of group 13 binary nitrides MN from organometallic precursors. Recently we proposed that carbon incorporation in MN is attributed to the metal-carbon bonding in the gas-phase with formation of organometallic heterocycles. To further explore processes of metal-carbon and metal-nitrogen bond formation, model complexes $X_3M\text{-NCR}$ and $X_3M\text{-CNR}$ ($M=\text{Al,Ga}$; $X=\text{H,Cl,CH}_3$; $R=\text{H,CH}_3$), metal cyanides $X_2\text{MCN}$ and $X_2\text{MNC}$ and their oligomers $[X_2\text{MCN}]_n$ ($n=2,3,4$) have been chosen for the present investigation. Geometries, relative energies, vibrational frequencies, and thermodynamic parameters of dissociation and elimination reactions are predicted at B3LYP level of theory with full-electron pVDZ basis set. All structures were fully optimized with subsequent vibrational analysis. Gaussian 94 set of programs was used throughout.

Results of this study indicate competitiveness of gallium-carbon and gallium-nitrogen bond formation. Predicted dissociation enthalpies of metal-carbon bonded complexes are about 15 kJ mol^{-1} *higher* compared to dissociation enthalpies of metal-nitrogen bonded isomers both for Al and Ga. For $X_2\text{MCN}$ species, Ga-C and Al-N bonding is found to be preferable compared to Ga-N and Al-C, in qualitative agreement with Pearson's HSAB concept. Relative stability of $X_2\text{MCN}$ forms is increasing with increasing of electronegativity of terminal group X both for Al and Ga: $\text{CH}_3 < \text{H} < \text{CN} < \text{Cl}$. RX elimination reactions with dimerization of $X_2\text{MCN}$ are extremely favorable in case of $X=\text{H,CH}_3$; in contrast, for $X=\text{Cl}$ dissociation processes are predominant.